

Available online on 15.12.2018 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

In vitro Clot Dissolving Activity of Carbomer Based Gel Containing Ethanolic Extract *Calotropis gigantea* Leaves

Prerana Alawa¹, Mayank Tenguria² and Vijayshri Nilosey¹¹ Department of Chemistry, Holkar Science College, Indore, MP, India² Lenience Biotech Lab, Bhopal, MP, India

ABSTRACT

Blood clotting is an important and vital process that takes places in humans, animals and birds. At the same time blood clots sometimes considered an issue when developed in the circulatory system due to failure of hemostasis causes vascular blockage and leads to serious consequences in thrombolytic diseases such as acute myocardial or cerebral infarction which may cause death. Certain substances like Alteplase, anistreplase, streptokinase, urokinase, tissue plasminogen activator, Heparin and Aspirin are used as clot dissolving agents. Plants have also been proved to perform such activity. *Calotropis gigantea* L. is a traditional medicinal plant with several pharmacological properties belongs to the family Apocyanaceae of *Asclepiadaceae* was investigated for its efficacy for blood clot dissolving activity under *in vitro* conditions when incorporated into carbomer based gels. The ethanolic extract of the leaves of *C. gigantea* was prepared by soxhlation which upon analysis reported to contain alkaloids, glycosides, flavonoids, tannins, terpenoids and saponins as usual components as were reported in earlier investigations. There were 3 gels with ethanolic leaf extract *C. gigantea* with 1 mg, 0.5 mg and 0.25 mg per gram of 1% Carbopol-940 gel containing methyl paraben, glycerol and water. The two gels containing sterile distilled water and 0.1 mg per gram gel streptokinase enzyme served as -ve and positive controls respectively. The *in vitro* thrombolytic of clot dissolving activity was performed on goat blood collected from slaughter houses. From the results the percentage clot dissolving activity of gels containing 1 mg, 0.5 mg and 0.25 mg *C. gigantea* extract per gram were reported as 34%, 27% and 14% respectively considered significant when compared to the activity of standard streptokinase in the gel. The activity of these experimental gels describes the possibility of development of new safe, and non reactive topical anti-clot gel products using herbal or plant components like *C. gigantea* extracts.

Keywords: *Calotropis gigantea* L, Streptokinase, Thrombolytic activity, Carbopol gel, Biopharmaceutical.**Article Info:** Received 18 Oct 2018; Review Completed 29 Nov 2018; Accepted 30 Nov 2018; Available online 15 Dec 2018

Cite this article as:

Alawa P, Tenguria M, Nilofey V, *In Vitro* Clot Dissolving Activity of Carbomer Based Gel Containing Ethanolic Extract of *Calotropis Gigantea* Leaves, Journal of Drug Delivery and Therapeutics. 2018; 8(6-s):101-104**DOI:** <http://dx.doi.org/10.22270/jddt.v8i6-s.2093>

*Address for Correspondence:

Mayank Tenguria, Lenience Biotech Lab, Bhopal, MP, India

INTRODUCTION

Referring to the article published in 1879, blood clotting is an important and vital process that takes places in humans, animals and birds¹. Maintenance of normal blood flow requires equilibrium between procoagulant and anticoagulant factors within the blood circulatory system. But when procoagulant activity preponderates, results into the formation of blood clots where tissue damage becomes a factor in the process in general². Blood clotting is a complex interaction of various mechanisms, where activation of the coagulation, fibrinolytic systems, disruption of the vascular endothelium, and the generalized activation of cellular mechanisms leads to clotting on the surface of monocytes and platelets in circulation³. Blood clots sometimes considered an issue when developed in the circulatory system due to failure of hemostasis causes vascular blockage and leads to serious consequences in thrombolytic diseases such as acute

myocardial or cerebral infarction which may cause death⁴. Hereditary factors, primary or acquired, play a role in the development of thrombosis³. The breakdown of blood clots is referred to as thrombolysis or clot busting in pharmacological terms and that can be done by stimulating secondary fibrinolysis by plasmin through infusion of analogs of tissue plasminogen activator the protein that normally activates plasmin. Certain substances are used as Clot dissolving agents available in the market like Alteplase, anistreplase, streptokinase, urokinase and tissue plasminogen activator. Where moderately efficient Heparin and Aspirin are also available accelerating the lysis and prevent the reoccurrence of thrombus but are safe⁵. As per some literatures, anticoagulants that are being used presently shows not only the negative and life-threatening side effects but runs high in cost from therapeutic point of view, thus researchers and investigators are now finds their interest in exploring the natural resources in order to develop alternate novel agent that could perform similar

functions without any harm ^{6,7}. According to World Health Organization (WHO), medicinal plants would be the best source to obtain variety of drugs, due to the presence of different chemicals used as phytomedicine, derived from different parts of the plants⁸. In recent studies, *Campomanesia xanthocarpa*, *Alpinia conchigera*, *Lannea grandis* Engl. *Aglaonema hookerianum* Schott. *Tridax procumbens* L. *Nigella sativa*, *Capsicum frutescens*, *Brassica oleracea*, *Calotropis gigantea* like plants have been proved to perform such activity⁹⁻¹². *Calotropis gigantea* L. is a traditional medicinal plant has reported to have analgesic, antipyretic activity, antimicrobial, anti-diarrheal and cytotoxic, properties belongs to the family Apocyanaceae of *Asclepiadaceae* thrives habitat of Asian countries¹³. *C. gigantea* L. has been considered to cure cardiovascular diseases, skin diseases, use as anticancer, antifertility and antidote for snakebites in traditional system of medicine¹⁴⁻¹⁷. This plant has also been investigated for thrombolytic activity¹². Thus current preliminary study aimed at investigation efficacy of carbomer based gel prepared from ethanolic extracts of *C. gigantea* leaves blood clot dissolving activity under *in vitro* conditions.

MATERIALS AND METHODS

Sample collection and processing

The leaves of *C. gigantea* were collected from local area around the Indore City and were authenticated with use of herbarium available in the college, literatures and images from web resources. The collected leaves were washed thoroughly with tap water, cleaned and dried at room temperature followed by grinding them into fine powder. These finely grounded leaf powders were subjected to phytochemical extraction.

Phytochemical extraction and analysis

The fine powdered leaves of *C. gigantea* were first defatted with petroleum ether at ambient temperature for 24 hours and then the defatted marc was further subjected to phytochemical extraction using Soxhlet extractor in 50% ethanol as solvent. The menstruum so obtained after extraction was allowed to concentrate in a boiling water bath to evaporate the solvent. The percentage yield of extraction and preliminary phytochemical analysis was done¹⁸⁻²⁰.

Preparation of carbopol based gel

Stock of phytochemical extract was prepared in sterile distilled water with a concentration of 100 mg/ml. In present study the Carbopol gel were prepared with suitable modifications ²¹⁻²³. The composition of the gel base used in present work is depicted in table 1. The stock concentration of 100 mg/ml crud phytochemical extract was prepared in sterile distilled water homogenized and degassed in ultrasonic bath which was used to be added in gel to make varied final concentrations of extract in per gram of gel.

Table 1 Composition of Cabopol-940 gel base

S.N.	Composition	Amount
1	Water	100 ml
2	Carbopol -940	1 gm
3	Methyl paraben	200 mg
4	Glycerol	1 ml
5	Phyto extract stock	As required

In vitro clot dissolving activity

The *in vitro* thrombolytic activity of prepared gels was done with some modifications^{5, 24, 25}. The blood samples were collected from the slaughter houses where goat blood was used instead of human blood because of similarities in clotting properties. The blood samples were collected in 1.5 ml microfuge and incubated at 37°C for 2 hours to allow the clot formation followed by removal of serum carefully and weight of the clot was measured. Now, to each of the pre-weighed clot in microfuge, 100 µl of prepared carbopol based gel containing extracts were added with the help of micropipette (from HiMedia) touching the surface of blood clot. A blank gel without any extract and a gel containing streptokinase were used as negative and positive control respectively. The set of experiment was incubated at 37 °C for 2 hours followed by observation *in vitro* thrombolytic activity. The liquid released after incubation was drained and remaining clot was washed with sterile distilled water and dried well to remove moisture followed by weighing the clot again to determine the percentage thrombolytic activity of gels compared to standard using following formulae;

$$\% \text{ Thrombolysis} = \frac{\text{Weight of clot after incubation}}{\text{Weight of clot before incubation}} \times 100$$

RESULTS AND DISCUSSION

Preliminary phytochemical analysis

The results of phytochemical extraction and analysis for the presence of phytoconstituents are more or less coincides with earlier investigations²⁶⁻²⁸. In present study the yield of extraction with hydro ethanolic solvent was 18.1%. While this ethanolic extract of leave of *C. gigantea* was reported to contain alkaloids, glycosides, flavonoids, tannins, terpenoids and saponins as usual components as were reported in earlier investigations²⁹.

In vitro clot dissolving activity

The *in vitro* clot dissolving activity of carbopol based gels shows promising results in development of biopharmaceutical purpose for therapeutic substances. There were 3 different concentrations of extracts in gels were used in present study whose percentage thrombolytic activity are depicted in table 2 compared to the standard streptokinase.

Table 2: *In vitro* clot dissolving activity of carbopol gel containing *C. gigantea* extract compared to gel containing Streptokinase

S.N.	Gels	Designation	Concentration	Percentage Clot Lysis
1	Gel-1	-ve Control with water	NA	1.8%
2	Gel-2	+ve Control with Streptokinase	0.1 mg/g gel	71%
3	Gel-3	Test Gel with extract	1 mg/g gel	34%
4	Gel-4	Test Gel with extract	0.5 mg/g gel	27%
5	Gel-5	Test Gel with extract	0.25 mg/g gel	14%

The early reports on *in vitro* thrombolytic activity of *C. gigantea* extracts encouraged the test the similar effect of ethanolic extracts of *C. gigantea* leave when incorporated in any neutral gels in present study³⁰⁻³¹. In present study the *in vitro* blood clot dissolving activity of ethanolic leaf extract of *C. gigantea* in carbopol gel at a concentration of 1 mg/g of gel was observed to be 34% which reduces with serial decrease in extract concentration in gels as 27% and 14% with Gel-4 and Gel-5 respectively (Table 2). Streptokinase at a concentration of 0.1 mg/ml generally produces a thrombolytic activity in range from 80 to 95%^{12, 32}. In present study when it was incorporated in gel it showed an activity of 71% at similar concentration in gels that is optimum. The activity of varied concentration of extract with the activity streptokinase in gels, the dissolution of clot by extracts is considerable (Figure 1)




Figure 1: Graphical representation of *in vitro* thrombolytic or blood clot dissolving potential of gels prepares containing streptokinase and ethanolic extract of *C. gigantea*.

Food and Drug Administration (FDA) has approved several pharmaceuticals of plant origin, also several plants have been investigated earlier including *C. gigantea* for anti-coagulant, anti-platelet and fibrinolytic activity where many claims that some herbs and food showing preventive effects on events of coronary blockage and stroke^{33,34}. Heavy external blood clots due to any tissue tear or rupture or large accidental wound development is also a problem sometimes when causing painful bandage. Incorporation of any pharmacological substance in gel are generally meant for topical applications and here the anti blood clot activity of carbopol based gel containing ethanolic extracts of *C. gigantea* leaves were effectively dissolving blood clot at all 3 different concentration under *in vitro* experiments. This property of *C. gigantea* leaf extract may be exploited in development of new biopharmaceutical and therapeutic agents upon further extensive investigation related to it are save use on animals and humans.

CONCLUSIONS

From the results of present investigation, it could be concluded that the phytochemical extracts of *C. gigantea* are rich in number of chemical components that posses the biological and pharmacological activities which has been also reported several times in early studies. The clot dissolving activity of extracts works when not only when use as sole component but also when incorporated in gels. Because of the significant inadequacies of the currently available thrombolytic agents of microbial origin sometimes imparting allergy effects where plant derived drugs are considered promising in development of new therapeutic agents. So, upon further extensive studies related to preparation of non reactive polymer based gels containing plant chemicals could be developed into a successful topical biopharmaceutical product with anti-clot activity.

REFERENCES

- Barnes R, The Significance of thrombosis or blood clotting, The British Medical Journal, 1879; 1:966-968.
- Rubio-Jurado B, Iniguez-Franco P, Reyes PA, Robles G, Salazar-Páramo M, Zavala-Cerna G, et al., The clinical significance of coagulation and the inflammatory response in autoimmunity, Clinical Reviews in Allergy & Immunology, 2012; 42(2):172-80.
- Rickles FR, Falanga A, Molecular basis for the relationship between thrombosis and cancer, Thrombosis Research, 2001; 102:215-224.
- Sherwani SK, Bashir A, Haider SS, Shah MA, Kazmi SU, Thrombolytic potential of aqueous and methanolic crude extracts of camellia sinensis (green tea): *in vitro* study, Journal of Pharmacognosy and Phytochemistry, 2013; 2(1):125-129.
- Fatema T, Somaia HC, Kamrun NM, Imrul Hasif KM, Ahmed T, Akter M, *In-vitro* thrombolytic activity and phytochemical evaluation of leaf extracts of four medicinal plants of Asteraceae family, Journal of Pharmacognosy and Phytochemistry, 2017; 6(4):1166-1169.
- Manicam C, Abdullah JO, Mohd Tohit ER, Seman Z, Chin S, Hamid M, *In vitro* anticoagulant activities of Melastoma malabathricum Linn aqueous leaf extract: A preliminary novel finding, Journal of Medicinal Plants Research, 2010; 4(14):1464-1472.
- Kumar S, Joseph L, George M, Sharma A, A review on anticoagulant/antithrombotic activity of natural plants used in traditional medicine, International Journal of Pharmaceutical Sciences Review and Research, 2011; 8(1):70-74.
- Tenguria M, Ahirwar KK, Joshi PD, Estimation of total polyphenolic content and antibiogram studies of leaf and fruit aqueous extract of xanthium strumarium l, Science Secure Journal of Biotechnology, 2013; 2(3):83-88.
- Anwar MS, Khan IN, Barua S, Kamal ATMM, Hosen SMZ, Kawsar MH, Assessment of thrombolytic & cytotoxic activity of herbal preparations originated from botanical source of Bangladesh, Journal of Applied Pharmaceutical Science, 2011; 1(7):77-80.
- Klafke JZ, Da Silva MR, Rossato MF, Trevisan G, Walker CIB, Leal CAM, et al., Antiplatelet, antithrombotic and fibrinolytic activities of campomanesia xanthocarpa, Evidence-Based Complementary and Alternative Medicine, 2012; 8.
- Sultana I, Noor M, Barua J, Mahmood A, Das MC, Islam M, et al., *In-vitro* anti-atherothrombosis activity of four bangladeshi plants, International Journal of Green Pharmacy, 2012; 6:5-8.
- Ashrafudoulla M, Luna NA, Jyothy FH, Khatun T, Rafa TF, Parvin S, et al., Phytochemical and pharmacological investigation of calotropis gigantea, Journal of Medicinal Plants Studies, 2016; 4(4):245-255.
- Manivannan R, Shopna R, Anti-microbial and anti-inflammatory activity of new 4-methoxy-3-(methoxymethyl) phenol and (e)-n'-(5-bromo-2-methoxybenzylidene)-4-methoxy benzohydrazide isolated from calotropis gigantea white, Natural Product Sciences, 2017; 23(1):69-74.
- Suresh Babu AR, Karki SS, Wound healing activity of calotropis gigantea leaves in albino wistar rats, International Journal of Pharmacy, 2012; 2(1): 195-199.
- Park G, Lee EJ, Min HY, Choi HY, Han Ah-R, Lee SK, et al., Evaluation of cytotoxic potential of indonesian medicinal plants in cultured human cancer cells, Natural Product Sciences, 2002; 8(4):165-169.

16. Upendra B, Norton SP, Chandrakanth B, Role of uterine mast cells in developing anti-fertility effect of *Calotropis gigantea* Linn. in combination with ovarian hormones, *Indian Journal of Applied and Pure Biology*, 1992; 7(1):9-13.
17. Kitagawa I, Ru-song Z, Jony DP, Nam IB, Yasuyuki T, Mayasuki Y, et al., Chemical structure of calotropisides a and b two new oxypregnaneoligoglycosides from the root of *Calotropis gigantea* (Asclepiadaceae), *Chemical and Pharmaceutical Bulletin*, 1992; 40:2007-2013.
18. Harborne JB, *Phytochemical methods; a guide to modern techniques of plant analysis*, 3rd edition, Chapman and Hall, New York, 1983.
19. Khandelwal KR, *Practical pharmacognosy, technique and experiments*, 23rd edition, New Delhi, 2005.
20. Tenguria M, Chand P, Upadhyay R, Estimation of total polyphenolic content in aqueous and methanolic extracts from the bark of *Acacia nilotica*, *International Journal of Pharmaceutical Science and Research*, 2012; 3(9): 3458-3461.
21. Mohammad Haneefa KP, Shahima Hanan K, Sraswathi R, Mohanta GP, Nayar C, Formulation and evaluation of herbal gel of *Pothos scandens* Linn, *Asian Pacific Journal of Tropical Medicine*, 2016; 3(12):988-992.
22. Muhammad U, Ilyas, S, Mir S, Khan AK, Rashid R, Khan MZU, et al., Formulation and in vitro evaluation of carbapol 934 based modified clotrimazole gel for topical application, *Anais da Academia Brasileira de Ciencias*, 2016; 88(4):2303-2317.
23. Krongrawa W, Limmatvapirat S, Pongnimitprasert N, Meentam P, Limmatvapirat C, Formulation and evaluation of gels containing coconut kernel extract for topical application, *Asian Journal of Pharmaceutical Sciences*, 2018; 1-10.
24. Prasad S, Kashyap RS, Deopujari JY, Purohit HJ, Taori GM, Dagainawala HF, Development of an in vitro model to study clot lysis activity of thrombolytic drugs, *Thrombosis Journal*, 2006; 4:14.
25. Sweta P, Rajpal SK, Jayant YD, Hemant JP, Girdhar MT, Hatim FD, Effect of *Fagonia Arabica* (Dhamasa) on in vitro thrombolysis, *BMC Complementary and Alternative Medicine* 2007; 7: 36.
26. Singh S, Singh S, Mishra RM, Shrivastava MP, Preliminary phytochemical screening of *Calotropis gigantea* leaf, *International Journal of Scientific and Research Publications*, 2014; 4(2):1-3.
27. Radhakrishnan S, Alarfaj AA, Annadurai G, Estimation of phytochemical analysis and in vitro antioxidant activity of *Calotropis gigantea* extract: wound healing activity and its biomedical application, *International Journal of Pharmaceutical Science and Research*, 2015; 6(7):3053-3060.
28. Richa Harsimran K, Shikha S, Phytochemical and anatomical study of two species of *Calotropis* from Chandigarh, *International Journal of Pharmaceutical Science and Research*, 2015; 6(4):1452-1459.
29. Kori P, Alawa P, Evolution of anti microbial activity and phytochemical analysis of different organic solvent of *Calotropis gigantea*, *IOSR Journal of Pharmacy*, 2014; 4(7):50-55
30. Rajesh R, Raghavendra Gowda CD, Nataraju A, Dhananjaya BL, Kemparaju K, Vishwanath BS, Procoagulant activity of *Calotropis gigantea* latex associated with fibrin(ogen)olytic activity, *Toxicon*, 2005; 46:84-92.
31. Bindhu OS, Singh MK, Hemostatic, milk clotting and blood stain removal potential of cysteine proteases from *Calotropis gigantea* (L.) R. Br. Latex, *Pharmacognosy Magazine*, 2014; 10(Suppl 2):S350-S356.
32. Zaman R, Parvez M, Jakaria M, Sayeed MA, Islam M, In vitro clot lysis activity of different extracts of *Mangifera sylvatica* Roxb. Leaves, *Research Journal of Medicinal Plant*, 2015; 9(3):135-140.
33. Emran TB, Rahman MA, Uddin MMN, Rahman MMR, Uddin MZ, Dash R, et al., Effects of organic extracts and their different fractions of five bangladeshi plants on in vitro thrombolysis, *BMC Complementary Alternative Medicine* 2015; 15.
34. Ramjan AM, Hossain JF, Runa M, Hasanuzzaman, Mahmudul I, Evaluation of thrombolytic potential of three medicinal plants available in Bangladesh, as a potent source of thrombolytic compounds, *Avicenna Journal of Phytomedicine*, 2014; 4:430-436.

